

Renormalization in quantum Brans-Dicke gravity

Z. Haba

Institute of Theoretical Physics, University of Wrocław,
50-204 Wrocław, Plac Maxa Borna 9, Poland
e-mail: zhab@ift.uni.wroc.pl

Abstract

In the Brans-Dicke model we treat the scalar field exactly and expand the gravitational field in a power series. A comparison with 2D sigma models and ϕ^4 perturbation theory in four dimensions suggests that the perturbation series in 4D Brans-Dicke model is renormalizable.

1 Introduction

The appearance of a dimensional coupling constant in Einstein gravity supplies a simple argument of non-renormalizability of Einstein quantum gravity. The Brans-Dicke model [1] with the action

$$W = \int dx \left(\frac{1}{2\xi} \sqrt{g} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} \sqrt{g} \phi^2 R \right) \quad (1)$$

has been constructed in such a way that a scalar field appears in the place of the coupling constant. In such a case there is no dimensional coupling constant in the model. Unfortunately, in the Brans-Dicke Lagrangian (1) there is no free Lagrangian (quadratic in the metric tensor) which could be a starting point for a perturbative expansion. Our basic idea is to do the functional integration with an action which depends on the quantum scalar field but is quadratic in the metric field.

Another approach is discussed in [2]-[3]. By a conformal rescaling of the metric tensor in the classical action (1) $g_{\mu\nu} \rightarrow \phi^{-2} g_{\mu\nu}$ we have $\sqrt{g} \phi^2 R \rightarrow \sqrt{g} R$. Then, the authors can perform the perturbation expansion around a free action which is quadratic in the scalar field and (separately) in the metric field. It comes out to be non-renormalizable. However, the latter approach is not equivalent to the first one. The conformal (Weyl) invariance is lost at the quantum level because of the quantum anomalies. As a consequence Brans and Dicke's main idea of dimensionless coupling is lost in such an approach. Let us note that the loop expansion discussed in [2]-[3] when applied to the Brans-Dicke model

cannot be considered as the saddle point method (where $\hbar$ is to be the small parameter). In $W/\hbar$ in eq.(1) the parameter $\hbar$ can be absorbed into ϕ and there remains no small parameter which could serve for an expansion in both the metric and the scalar fields (we show however that our perturbation expansion in the metric field can be considered as an expansion in $\xi^{-\delta}$ with a certain $\delta > 0$). Let us note that there is some similarity between our functional integration over ϕ and a preliminary integration over longitudinal modes of the metric advocated in [4] -[5].

The Brans-Dicke model appears as the 1-loop approximation to the string theory [6][7][8]. In such a case the two approaches mentioned above are called the "string frame" and the "Einstein frame". From our point of view such a distinction is essential. This is the "string frame" applied at short distances which requires a renormalization. The "Einstein frame" whose quantum version is discussed in [2]-[3] applies to long distances and therefore can remain unrenormalized. It follows from our results that there is no need for the string tension parameter to serve as an ultraviolet cutoff for the effective string field theory. However, this dimensional parameter can distinguish the standard free behaviour of renormalizable quantum field theories from a regular behaviour on sub-planckian distances [9]. In a derivation of the effective string field theory only the massless modes are considered [10]. For this reason the model is scale invariant. The exact theory should contain mass parameters breaking the scale invariance.

The classical Brans-Dicke model of scalar-tensor gravity is scale invariant, i.e., $W(\tilde{g}, \tilde{\phi}) = W(g, \phi)$ with

$$\tilde{g}_{\mu\nu}(x) = g_{\mu\nu}(\lambda x) \quad (2)$$

and

$$\tilde{\phi}(x) = \lambda\phi(\lambda x)$$

In order to preserve positivity of the metric g we write

$$\sqrt{g}g^{\mu\nu} = (\exp h)^{\mu\nu} \equiv G^{\mu\nu} \quad (3)$$

where $h^{\mu\nu}$ is also dimensionless (in the classical model) and assumed to be small. Eq.(3) seems to be a reasonable starting point for an expansion around the flat space. However, in the conventional perturbation expansion of Einstein quantum gravity we would encounter a difficulty with the assumptions (2)-(3) because already at the lowest order h has a scale dimension 1. We show that this obstacle can be avoided in the Brans-Dicke model. We obtain a perturbation expansion which is consistent with the scaling (2).

We shall give arguments showing that the ultraviolet behaviour of the 4D Brans-Dicke theory is similar to the ultraviolet behaviour of (non-compact) σ -models in two dimensions. For this aim let us write down the action (1) in a way resembling an action of $GL(4, R)$ -valued sigma field [11]-[12]. So, we treat

G in eq.(3) as an element of $GL(4, R)$. We can represent the Christoffel symbol in the form (see ref.[11], Appendix A)

$$\Gamma = aJ + bG^{-1}JG \quad (4)$$

where

$$J_\mu = G^{-1}\partial_\mu G$$

Here, J is 4×4 matrix; in eq.(4) the components of Γ are expressed by matrix elements of J and G in a way which is irrelevant at the moment. Now, in $W(G) = \frac{1}{2} \int dx \phi^2 G^{\mu\nu} R_{\mu\nu}$ the Ricci tensor is of the form $\partial\Gamma - \Gamma\Gamma$. Hence, the final formula for the gravitational action $W(G)$ has the form resembling the σ -model

$$W(G) = \int dx \phi^2 G(a_1 J J + a_2 \partial J) \quad (5)$$

where a_k are certain constants. The precise contraction of indices is rather complicated and irrelevant for our discussion. The matrix notation is useful when we wish to follow the perturbation expansion. So, the lowest order in the expansion of $W(G)$ in terms of h is $W_1(G) = \int \phi^2 \partial^2 h$. Then, there is the bilinear term $\frac{1}{2} \phi^2 h \partial^2 h$ which will be considered as the free action $W_2(G)$. Now, an expansion of the interaction $W_I \equiv W(G) - W_1(G) - W_2(G)$ starts as follows

$$W_I(G) = \int dx \phi^2 (b_1 h \partial h \partial h + b_2 h h \partial^2 h + b_3 h h \partial h \partial h + b_4 h h h \partial^2 h + \dots) \quad (6)$$

with certain constants b_k .

2 The lowest order

We work in a generalization of the harmonic gauge

$$\partial_\mu(\phi^2 G^{\mu\nu}) = 0 \quad (7)$$

This gauge has the advantage that the bilinear form quadratic in the second order derivatives has an exceptionally simple form. So, the gravitational part $W(G)$ of the action (1) in the gauge (7) can be written as

$$W(G) = \frac{1}{4} \int dx (\phi^2_{,\nu} G^{\mu\nu} \Gamma_{\mu\sigma}^\sigma - \phi^2_{,\sigma} G^{\mu\nu} \Gamma_{\mu\nu}^\sigma + \phi^2 G^{\mu\nu} \Gamma_{\mu\nu,\sigma}^\sigma) \quad (8)$$

Then, the quadratic form which is of the second order in derivatives reads

$$W_2^Q(G) = -\frac{1}{8} \int dx \phi^2 \hat{h}^{\alpha\beta} \Delta \hat{h}^{\alpha\beta} + \frac{1}{32} \int dx \phi^2 H \Delta H \quad (9)$$

where Δ is the Laplace operator in R^4 ,

$$\hat{h}^{\alpha\beta} = h^{\alpha\beta} - \delta^{\alpha\beta} \frac{1}{4} H$$

is the traceless part of h and $H = Trh = h^{\alpha\alpha}$ denotes the trace of h . Then, the bilinear form resulting from W together with the Faddeev-Popov determinant for the gauge (7) define the integration measure

$$d\mu(h, \phi) = \mathcal{D}h \mathcal{D}\phi \exp(-W_2(h, \phi)) \det D_{FP}$$

where

$$W_2(h, \phi) = \frac{1}{2\xi} \int dx \phi(-\Delta)\phi + \frac{1}{4} \int dx \hat{h}^{\mu\nu}(-\Delta_{2T})\hat{h}^{\mu\nu} - \frac{1}{16} \int dx H(-\Delta_{2S})H \quad (10)$$

The Faddeev-Popov operator for the gauge (7) is

$$D_{FP} = \frac{1}{2} \phi^2 G^{\mu\nu} \partial_\mu \partial_\nu \quad (11)$$

The operators in the quadratic form in $W_2(G)$ can be deduced from eqs.(8)-(11). So,

$$\Delta_{2T} = \frac{1}{2} \phi^2 \Delta + \frac{1}{2} \phi_{,\nu}^2 C^{\nu\mu} \partial_\mu \quad (12)$$

Here, the operator Δ_{2T} is acting on tensors of the second order, the Laplacian Δ is considered as the diagonal operator in the space of tensors and $C^{\mu\nu}$ are matrices built from δ -functions whose explicit form can be read from eq.(8). The operator Δ_{2S} is acting on scalars and has the form

$$\Delta_{2S} = \frac{1}{2} \phi^2 \Delta - \frac{1}{4} \phi_{,\nu}^2 \partial_\nu \quad (13)$$

We can generalize the model (10) for an application of the dimensional regularization replacing the scalar free action $W_2(\phi) = \frac{1}{2\xi} \int dx \phi(-\Delta)\phi$ by

$$W_2^\gamma(\phi) = \frac{1}{2\xi} \int dx \phi(-\Delta)^{2-\gamma} \phi \quad (14)$$

The resulting Brans-Dicke action will be denoted W^γ . The action W^γ is also scale invariant $W^\gamma(\tilde{h}, \tilde{\phi}) = W^\gamma(h, \phi)$ where $\tilde{\phi}(x) = \lambda^\gamma \phi(\lambda x)$ and $\tilde{h}(x) = \lambda^{1-\gamma} h(\lambda x)$.

For a perturbation expansion instead of the determinant $\det D_{FP}$ the Faddeev-Popov ghosts $(\bar{\chi}, \chi)$ are introduced. The corresponding action is

$$W(\chi) = \frac{1}{2} \int dx \phi^2 G^{\mu\nu} \bar{\chi} \partial_\mu \partial_\nu \chi \quad (15)$$

It is scale invariant, i.e., invariant under the change of variables $\tilde{\chi}(x) = \lambda^{1-\gamma} \chi(\lambda x)$ (together with the scale change of ϕ and h mentioned before).

We apply some methods of our earlier paper [9] where an interaction with a scale invariant metric has been discussed. First, let us consider the propagator

for the Faddeev-Popov ghosts (15) in the proper time representation. We take an average over ϕ of the heat kernel K_τ of the operator D_{FP} . The path integral formula takes the form

$$\langle K_\tau(x, y) \rangle = \langle (\exp \tau D_{FP})(x, y) \rangle = \int \mathcal{D}\phi \exp(-W_2(\phi)) \int \mathcal{D}q \exp(-\frac{1}{2} \int \phi^2 G^{\mu\nu} \frac{dq_\mu}{dt} \frac{dq_\nu}{dt}) \delta(q(0) - x) \delta(q(\tau) - y) \quad (16)$$

In order to obtain the behaviour of $\langle K_\tau(x, y) \rangle$ for a small τ it is sufficient to assume the scale invariance of ϕ

$$\int dx \mathcal{L}(\lambda^\gamma \phi(\lambda x)) = \int dx \mathcal{L}(\phi(x)) = W_2(\phi) \quad (17)$$

In fact, introducing in eq.(16) a new functional integration variable $\tilde{\phi} = \lambda^\gamma \phi(\lambda x)$ and a new path $\tilde{q}$ defined on an interval $[0, 1]$ by

$$q(\tau s) - x = \tau^\sigma \sqrt{\xi}(\tilde{q}(s) - x) \quad (18)$$

where

$$\sigma = \frac{1}{2}(1 + \gamma)^{-1}$$

we can explicitly extract the τ -dependence of $\langle K_\tau(x, y) \rangle$

$$\begin{aligned} \langle K_\tau(x, y) \rangle &= \langle E[\delta(y - x - \tau^{\frac{1}{2}-\sigma\gamma} \sqrt{\xi} \tilde{q}(1))] \rangle \\ &= \xi^{-2} \tau^{-\frac{2}{1+\gamma}} F\left(\tau^{-\frac{1}{2(1+\gamma)}} \xi^{-\frac{1}{2}} (y - x)\right) \end{aligned} \quad (19)$$

where $E[.]$ denotes an expectation value over the paths $\tilde{q}$ and F is the probability distribution of $\tilde{q}(1)$. Then, an average of the propagator (in the proper time representation) behaves as

$$\begin{aligned} \langle (-D_{FP})^{-1}(x, y) \rangle &= \int_0^\infty d\tau \langle K_\tau(x, y) \rangle \\ &= \xi^{-1-\gamma} \int_0^\infty d\tau \tau^{-\frac{2}{1+\gamma}} F\left(\tau^{-\frac{1}{2(1+\gamma)}} (y - x)\right) \simeq \xi^{-1-\gamma} |x - y|^{-2+2\gamma} \end{aligned} \quad (20)$$

if $\gamma < 1$. For $\gamma = 1$

$$\langle (-D_{FP})^{-1}(x, y) \rangle \simeq -\xi^{-2} \ln(x - y)^2 \quad (21)$$

We can represent the Faddeev-Popov determinant by its heat kernel

$$\det D_{FP} = \exp \text{Tr} \ln D_{FP} = \exp \left(- \int_0^\infty \frac{d\tau}{\tau} \int dx (\exp(\tau D_{FP}) - 1)(x, x) \right) \quad (22)$$

It follows from the small τ behaviour (19) of the heat kernel (16) that the integral over the proper time in eq.(22) is finite. Hence, no renormalization is needed for the Faddeev-Popov determinant.

We consider next the heat kernel K_τ and the propagator of the operator Δ_{2T} . The path integral formula takes the form

$$\begin{aligned} \langle (-\Delta_{2T})^{-1}(x, y) \rangle &= \int_0^\infty d\tau \langle K_\tau(x, y) \rangle = \int_0^\infty d\tau \langle (\exp \tau \Delta_{2T})(x, y) \rangle \\ &= \int_0^\infty d\tau \int \mathcal{D}\phi \exp(-W_2(\phi)) \int \mathcal{D}q \exp(-\frac{1}{2} \int \phi^2 \frac{dq_\mu}{dt} \frac{dq^\mu}{dt}) T_\tau \delta(q(0) - x) \delta(q(\tau) - y) \end{aligned} \quad (23)$$

where the matrix T_τ is the solution of the equation

$$dT_\tau = \phi_{,\mu} C^{\mu\nu} T_\tau dq_\nu \quad (24)$$

and the stochastic integral in eq.(24) is of the Ito type (see ref. [13]). The formula (23) is analogous to the well-known Feynman formula for a particle in an electromagnetic field. We obtain a similar formula for the heat kernel of Δ_{2S} . The argument with the change of variables (17)-(18) applies well to the functional integral (23). The change of time in T_τ is irrelevant for the short time behaviour of the τ -integral in (23) because T_τ is continuous in τ . Hence, the argument used at the propagator for ghosts applies here as well leading to the result

$$\begin{aligned} \langle (-\Delta_{2T})^{-1}(x, y) \rangle &= \int_0^\infty d\tau \langle K_\tau(x, y) \rangle \\ &\simeq \xi^{-1-\gamma} |x - y|^{-2+2\gamma} \end{aligned} \quad (25)$$

if $\gamma < 1$ and for $\gamma = 1$ we obtain the behaviour (21). It is clear that also

$$\langle (-\Delta_{2S})^{-1}(x, y) \rangle \simeq \xi^{-1-\gamma} |x - y|^{-2+2\gamma}$$

Strictly speaking the Euclidean functional integral over H is divergent. It has to be interpreted by means of a contour rotation in the functional space of metrics, see [4][14].

3 A perturbation expansion

The functional integral can be expressed in the form

$$d\mu_0(\phi) d\nu_T(\hat{h}) d\nu_S(H) \det(-\Delta_{2T})^{-\frac{1}{2}} \det(-\Delta_{2S})^{-\frac{1}{2}} \det D_{FP} \quad (26)$$

where μ_0 is a Gaussian measure with the covariance $\xi(-\Delta)^{-1}(x, y)$, ν_T is a Gaussian measure with the covariance $(-\Delta_{2T})^{-1}(x, y)$ and ν_S is a Gaussian measure with the covariance $(-\Delta_{2S})^{-1}(x, y)$. The determinants of Δ_{2T} and Δ_{2S} require no renormalization by the same argument as the one used in the discussion of $\det D_{FP}$. Eq.(21) means that

$$\int d\mu_0(\phi) \int d\nu_T(\hat{h}) \hat{h}_{\mu\nu,\rho}(x) \hat{h}_{\alpha\beta,\sigma}(y) \simeq -\xi^{-2} \partial_\rho \partial_\sigma \ln(x - y)^2 \quad (27)$$

The higher order correlations are expectation values over ϕ of sums of products of $(-\Delta_2)^{-1}$

$$\begin{aligned} \langle \hat{h}(x_1) \dots \hat{h}(x_{2n}) \rangle &= \sum_{(i,j)} \langle \Pi_{(i,j)} (-\Delta_{2T})^{-1}(x_i, x_j) \rangle = \int_0^\infty d\tau_1 \dots d\tau_n \\ \langle E[T_{\tau_1} \delta \left(x_2 - x_1 - \tau_1^{\frac{1}{2} - \sigma\gamma} \sqrt{\xi} \tilde{q}_1(1) \right) \dots T_{\tau_n} \delta \left(x_{2n} - x_{2n-1} - \tau_n^{\frac{1}{2} - \sigma\gamma} \sqrt{\xi} \tilde{q}_n(1) \right)] \rangle \\ &+ \text{permut.} \end{aligned} \quad (28)$$

where on the r.h.s. we have a sum over permutations of x_k in accordance with the well-known Gaussian combinatorics. It can be shown that the singularity of an expectation value of a product of $(-\Delta_{2T})^{-1}$ is the same as that of a product of singularities of $\langle (-\Delta_{2T})^{-1} \rangle$. It is clear that when we are applying formally the propagator (16) for the h -average then an expansion in h has the same singularity as an expansion in the two-dimensional σ -model. The latter is known to be renormalizable. In the Brans-Dicke model the problem is more complicated because in the perturbation expansion we have the mixed ϕh terms which lead to additional singularities. However, it can be seen from eqs.(5)-(6) that the first term in eq.(5) contributes to the interaction W_I terms of the form $\psi^2 \phi^2$ times powers of a field h of zero dimension where $\psi = \partial h$ and ϕ are fields of dimension 1. Hence, by power counting the perturbation expansion has the same singularity as the one in the ϕ^4 model (or a $\phi^2 \psi^2$ model). The second term in eq.(5) gives in W_I terms of the form $\phi^2 B$ times fields of dimension zero where $B = \partial \partial h$ is a field of dimension 2. The singularity of such an interaction is of the same kind as the one coming from the $\phi^2 \psi^2$ interaction. From the part of the Lagrangian (1) with derivatives of the scalar field we obtain an interaction of the form $W_I(\phi) = \int dx (G - 1) \partial \phi \partial \phi$. A renormalization of the perturbation expansion of this interaction is equivalent to the well-studied problem of the renormalization of a composite field (the energy-momentum tensor) in the free massless scalar quantum field theory. We have shown in sec.2 that the Faddeev-Popov determinant requires no renormalization. However, if we expand the action (15) around $G = 1$ then we obtain the same problem of a renormalization of the composite field as the one concerning the scalar field.

Calculating the terms in the perturbation expansion explicitly we can see that there are quartically divergent vacuum diagrams, as for example (with certain powers n and m)

$$\int dx dy \langle \phi^2(x) h^n(x) \partial h(x) \partial h(x) \phi^2(y) h^m(y) \partial h(y) \partial h(y) \rangle$$

which are analogs of $\int dx dy \langle \phi^2(x) \psi^2(x) \phi^2(y) \psi^2(y) \rangle$ in the $\phi^2 \psi^2$ model.

We can check renormalizability of connected diagrams (and the correspondence to the $\phi^2 \psi^2$ model) at lower orders again by explicit calculations. So, let us consider the two-point function $\langle h(x_1) h(x_2) \rangle$ in the second order of W_I

$$\int dx dy \langle \phi^2(x) h^n(x) \partial h(x) \partial h(x) :: \phi^2(y) h^m(y) \partial h(y) \partial h(y) : h(x_1) h(x_2) \rangle \quad (29)$$

From $\langle \phi^2 \phi^2 \rangle$ we obtain $(y-x)^{-4}$ and from $\langle \partial h(x) \partial h(y) \rangle$ we have $(x-y)^{-2}$. So, the connected diagrams are quadratically divergent. This is an analog of the diagram

$$\langle : \phi^2(x) \psi^2(x) :: \phi^2(y) \psi^2(y) : \psi(x_1) \psi(x_2) \rangle$$

which leads to the ψ -mass renormalization in the $\phi^2 \psi^2$ model (this will be the wave function renormalization in the model (5)). The four-point function at the same order would be logarithmically divergent. The divergence could be cancelled by the coupling constant renormalization (of the $\phi^2 \psi^2$ interaction). Higher order connected correlation functions are finite at the second order of W_I .

We could continue with higher orders. So, the third order term in the interaction leads to the higher order wave function renormalization in the propagator

$$\int dy_1 dy_2 dy_3 dy_4 \langle \phi^2(y_1) h^n(y_1) \partial h(y_1) \partial h(y_1) \phi^2(y_2) h^m(y_2) \partial h(y_2) \partial h(y_2) \phi^2(y_3) h^r(y_3) \partial h(y_3) \partial h(y_3) h(x_1) h(x_2) \rangle \quad (30)$$

In general, we can see that the expansion in h can be considered as an expansion in powers of $\xi^{-\frac{1}{2}-\frac{\gamma}{2}}$.

A complete proof of the renormalizability still requires a more detailed analysis but we think that we have given here some convincing arguments showing that a functional integration over the scalar field substantially improves the ultraviolet behaviour of the perturbation series. Another interesting question concerns the scale invariance of the summed up renormalized series. Although at each order the metric field h has zero dimension the logarithms can sum up in $G = \exp h$ leading to a positive dimensionality of the metric field G . In fact, if we treat the matrix h like a commutative Gaussian field then

$$\int d\mu_0(\phi) \int d\nu_T(\hat{h}) : \exp \hat{h}(x) :: \exp \hat{h}(y) : \simeq \exp \frac{1}{2} \langle (-\Delta_{2T})^{-1}(x, y) \rangle \quad (31)$$

The scale dimension of the metric (if positive) determines the short distance behaviour of matter fields [9]. If the approximation (31) is reliable then a determination of the constants in eq.(21) would determine the short distance behaviour of all quantum fields interacting with gravity.

References

- [1] C. Brans and R.H. Dicke, Phys.Rev.**124**,925(1961)
- [2] S. Deser and P. van Nieuwenhuizen, Phys.Rev.**D10**,401(1974)
- [3] G. 't Hooft and M. Veltman, Ann.Inst.H. Poincare,**20A**,69(1974)
- [4] G.W. Gibbons, S.W. Hawking and M.J. Perry, Nucl.Phys.**B 138**,141(1978)
- [5] K. Schleich, Phys.Rev. **D36**,2342(1987)

- [6] A.A. Tseytlin, Phys. Lett.**B176**,92(1986)
- [7] C. G. Callan, C.Lovelace, C.R. Nappi and S.A. Yost,
Nucl. Phys.**B288**,525(1987)
- [8] G. Veneziano, Status of string cosmology in String Gravity and Physics at
the Planck Energy Scale, ed. by N. Sanchez and A. Zichichi,
Kluwer,Dordrecht,1996
- [9] Z. Haba,Phys.Lett.**B528**,129(2002)
- [10] C. G. Callan, I.R. Klebanov and M.I. Perry, Nucl. Phys.**B278**,78(1986)
- [11] D.M. Capper, G. Leibbrandt and M. Ramon Medrano,
Phys.Rev.**D8**,4320(1973)
- [12] A. Strominger, Phys. Rev.**D24**,3082(1981)
- [13] N. Ikeda and S. Watanabe, Stochastic Differential Equations and Diffusion
Processes, North-Holland, Amsterdam, 1981
- [14] P. O. Mazur and E. Mottola, Nucl.Phys.**B341**,187(1990)